VOLUME 58 • AUTHOR INDEX

Agabiti N *et al*. Short term respiratory effects of acute exposure to chlorine due to a swimming pool accident, **399**
Ahlgren T *see* Björk J *et al*
Ahn K-D *see* Todd AC *et al*
Albin M *see* Björk J *et al*
Aldrich JM *see* Kales SN *et al*
Alexopoulos EC, Burdorf A. Prognostic factors for respiratory sickness absence and return to work among blue collar workers and office personnel, **246**
Alfredsson L *see* Bildt C *et al*
Almeida JWR *see* Faria MAM and Almeida JWR
Ancona C *see* Agabiti N *et al*
Anderson HR *et al*. Particulate matter and daily mortality and hospital admissions in the west midlands conurbation of the United Kingdom: associations with fine and coarse particles, black smoke and sulphate, **504**
Andersson E *et al*. A case-referent study of cancer mortality among sulfate mill workers in Sweden, **321**
Andersson M *see* Iregren A *et al*
Anees W *et al*. Glutaraldehyde induced asthma in endoscopy nursing staff: *letter*, **544**
Anger WK *see* Engel LS *et al*
Angerer J *see* Kraus T *et al*
Ariëns GAM *et al*. Are neck flexion, neck rotation, and sitting at work risk factors for neck pain? Results of a prospective cohort study, **200**
Armstrong B *see* Sebastián MS *et al*
Aru G *see* Carta P *et al*
Atkinson RW *see* Anderson HR *et al*
Attfield MD *see* Chen W *et al*
Auburtin G *see* Cowie HA *et al*
Aw T-C *see* Bradberry SM *et al*
see Sadhra S *et al*
Axelsson G *see* Wennborg H *et al*

Bakke B *see* Ulvestad B *et al*
Ball LM *see* Schulz MR *et al*
Baris I *see* Dumortier P *et al*
Bartlett T *see* Goodfellow FJL *et al*
Bauer TT *et al*. Functional variables associated with the clinical grade of dyspnoea in coal miners with pneumoconiosis and mild bronchial obstruction, **794**
Baxter D *see* Cherry N *et al*
see Cherry N *et al*
Beach JR *see* Sadhra S *et al*
Beard KK *see* Burns CJ *et al*
Becher G *see* Jaakkola JJK *et al*
Beck J *see* Cowie HA *et al*
Beck W *see* Seidler A *et al*
Beral V *see* Doyle P *et al*
Berg NP *see* Seldén AI *et al*
Berglind N *see* Rosenlund M *et al*
Bertrand J-P *see* Nadif R *et al*
Best N *see* Keegan T *et al*
Bickeböller R *see* Seidler A *et al*
Bildt C *et al*. Effects of drop out in a longitudinal study of musculoskeletal disorders, **194**
Billström R *see* Björk J *et al*
Binotti M *see* Palin LA *et al*
Björk J *et al*. Are occupational, hobby, or lifestyle exposures associated with Philadelphia chromosome positive chronic myeloid leukaemia?, **722**
Blain PG. Adverse health effects after low level exposure to organophosphates: *editorial*, **689**
Bluhm G *see* Rosenlund M *et al*
Bodin L *see* Wennborg H *et al*
Bodin LS *see* Seldén AI *et al*
Boffetta P *et al*. Mortality from cardiovascular diseases and exposure to inorganic mercury, **461**
Bolm-Audorff U *see* Seidler A *et al*
Bona G *see* Palin LA *et al*
Bongers PM *see* Ariëns GAM *et al*
Book reviews Brimblecombe P, Maynard R, editors. The urban atmosphere and its effects., **683**
Fairman R, Murray V, Kirkwood A, *et al*, *editors*. Chemical incident management: local authority environmental health practitioners, **753**
Gardner DE, Crapo JD, McClellan RO, *editors*. Toxicology of the lung: 3rd edition, **545**
Harris RL, *editor*. Patty's industrial hygiene, Volumes 1–4, 5th edition., **684**
Health and Safety Executive, *editor*. The successful management of hand-arm vibration: interactive CD-ROM package, **280**
Hinds WC; Aerosol technology, second edition, **65**
Lewis RJ, *author*. Rapid guide to hazardous chemicals in the workplace, 4th edition., **65**
Seaton A, Seaton D, Leitch AG, *editors*. Crofton and Douglas's respiratory diseases, 5th edition., **137**
Williams PL, James RC, Roberts SM, *editors*. Principles of toxicology: environmental and industrial applications, 2nd edition., **545**
Wood M, Smith MW, *editors*. Health and safety in laboratory animal facilities., **280**
Bourke JF *see* O'Connor TM *et al*
Bouter LM *see* Ariëns GAM *et al*
Bovenzi M *et al*. Response of finger circulation to energy equivalent combinations of magnitude and duration of vibration, **185**
Bradberry SM *et al*. Occupational methaemoglobinaemia, **611**
Bremner SA *see* Anderson HR *et al*
Brennan N *see* O'Connor TM *et al*
Brenner H *see* Siebert U *et al*
Brodkin CA *et al*. Serum hepatic biochemical activity in two populations of workers exposed to styrene, **95**
Broucke I *see* Dumortier P *et al*
Buchanan D *see* Pilkington A *et al*
Buchanan D *et al*. Estimation of cumulative exposure to organophosphate sheep dips in a study of chronic neurological health effects among United Kingdom sheep dippers, **694**
Bulbulyan M *see* Boffetta P *et al*
Burdorf A *see* Alexopoulos EC and Burdorf A
see Elders LAM and Burdorf A
Burge PS *see* Anees W *et al*
see Chaiear N *et al*
Burns CJ *see* Freedman M and Burns CJ
Burns CJ *et al*. Mortality in chemical workers potentially exposed to 2,4-dichlorophenoxyacetic acid (2,4-D) 1945–94: an update, **24**

Caballero J-D *see* Boffetta P *et al*
Calvert G *see* Steenland K *et al*
Camicioli R *see* Engel LS *et al*
Camp J *see* Brodkin CA *et al*
Carere A *see* Crebelli R *et al*
Carpenter M *see* Egedahl R *et al*
Carrivick PJW *et al*. Consultative team to assess manual handling and reduce the risk of occupational injury, **339**
Carta P *et al*. Mortality from lung cancer among silicotic patients in Sardinia: an update study with 10 more years of follow up, **786**
Cartmill JB *see* Burns CJ *et al*
Ceccarelli F *see* Boffetta P *et al*
Chaiear N *et al*. Sensitisation to natural rubber latex: an epidemiological study of workers exposed during tapping and glove manufacture in Thailand, **386**
Checkoway H *see* Brodkin CA *et al*
see Engel LS *et al*
see Rice FL *et al*
Chen BT *see* Chen W *et al*
Chen J-Q *see* Chen W *et al*
Chen R *et al*. Environmental tobacco smoke and lung function in employees who never smoked: the Scottish MONICA study, **563**
Exposure to organic solvents and personality, **14**
Nasal, eye, and skin irritation in dockyard painters, **542**
Chen W *et al*. Exposure to silica and silicosis among tin miners in China: exposure-response analyses and risk assessment, **31**
Cherrie B *see* Buchanan D *et al*
Cherrie JW *see* Cowie HA *et al*
see Dennekamp M *et al*
see Dick CAJ *et al*
Cherry NM *et al*. Health and exposures of United Kingdom Gulf war veterans. Part I: The pattern and extent of ill health, **291**
Health and exposures of United Kingdom Gulf war veterans. Part II: The relation of health to exposure, **299**
Occupational exposure to solvents and male infertility, **635**
Cherry NM *see* Nahit ES *et al*
Choi SY *see* Park JW *et al*
Christiani DC *see* Kales SN *et al*
Clarke M *see* Kurinczuk JJ and Clarke M
Clouter A *see* Donaldson K *et al*
Cockcroft A. Continuing Professional Development in *Occupational and Environmental Medicine*: *editorial*, **1**
Getting *Occupational and Environmental Medicine* to developing countries: *editorial*, **425**
Coggon D *see* Noël B *et al*
Coggon D. Ethics in occupational health research, **685**
Monitoring trends in occupational illness: *editorial*, **691**
Cole DC *et al*. Work correlates of back problems and activity restriction due to musculoskeletal disorders in the Canadian national population health survey (NPHS) 1994–5 data, **728**
Collins J *see* Cherry N *et al*
Cooper C *see* Noël B *et al*
Çoplü L *see* Dumortier P *et al*
Corbo GM *see* Agabiti N *et al*
Córdoba JA *see* Sebastián MS *et al*

Corrias GA *see* Mocci F *et al*
Courage C *see* Duarte-Davidson R *et al*
Cowie HA *et al*. An epidemiological study of the respiratory health of workers in the European refractory ceramic fibre industry, **800**
Craig DS *see* Whyte P *et al*
Crebelli R *et al*. Exposure to benzene in urban workers: environmental and biological monitoring of traffic police in Rome, **165**
Creed F *see* Cherry N *et al*
see Cherry N *et al*
Cullinan P *see* Chaiear N *et al*
Currò F *see* Fabiano B *et al*

Daniel U *see* Siebert U *et al*
Danuser B *see* Radon K *et al*
Davies G *see* Doyle P *et al*
Dawson D *see* Reid K and Dawson D
de Boeck M *see* Lison D *et al*
de Maertelaer V *see* Dumortier P *et al*
Dennekamp M *see* Dick CAJ *et al*
Dennekamp M *et al*. Ultrafine particles and nitrogen oxides generated by gas and electric cooking, **511**
de Vuyst P *see* Dumortier P *et al*
Dick CAJ *see* Dennekamp M *et al*
Dick CAJ *et al*. Stimulation of IL-8 release from epithelial cells by gas cooker PM_{10}: a pilot study, **208**
Dick F *see* Chen R *et al*
see Chen R *et al*
Diderichsen F *see* Voss M *et al*
Dietz MC *see* Ihrig A *et al*
Di Napoli A *see* Agabiti N *et al*
Dodds L, King WD. Relation between trihalomethane compounds and birth defects, **443**
Doll SR. Defamatory article by Martin Walker: *correspondence*, **136**
Donaldson K *see* Dennekamp M *et al*
see Dick CAJ *et al*
Donaldson K *et al*. Ultrafine particles, **211**
Doolette DJ *see* Whyte P *et al*
D'Orsi F *see* Agabiti N *et al*
Douwes J *see* Eduard W *et al*
Douwes M *see* Ariëns GAM *et al*
Doyle P *et al*. Primary infertility in nuclear industry employees: report from the nuclear industry family study, **535**
Duarte-Davidson R *et al*. Benzene in the environment: an assessment of the potential risks to the health of the population: *review*, **2**
Dumortier P *et al*. Erionite bodies and fibres in bronchoalveolar lavage fluid (BALF) of residents from Tuzköy, Cappadocia, Turkey, **261**
Dunn G *see* Cherry N *et al*
see Cherry N *et al*
Dybing E *see* Jaakkola JJK *et al*

Echeverria D *see* Brodkin CA *et al*
Eduard W *see* Ulvestad B *et al*
Eduard W *et al*. Short term exposure to airborne microbial agents during farm work: exposure-response relations with eye and respiratory symptoms, **113**
Egedahl R *et al*. Mortality experience among employees at a hydrometallurgical nickel refinery and fertiliser complex in Fort Saskatchewan, Alberta (1954–95), **711**
Ehrenberg L *see* Granath F *et al*
Eibel R *see* Bauer TT *et al*
Elders LAM, Burdorf A. Interrelations of risk factors and low back pain in scaffolders, **597**
Elliott P *see* Keegan T *et al*
Elon L *see* Williams PL *et al*
Elovainio M *see* Kivimäki M *et al*
Elsner G *see* Seidler A *et al*
Emri S *see* Dumortier P *et al*
Engel LS *et al*. Parkinsonism and occupational exposure to pesticides, **582**
Eyles J *see* Cole DC *et al*

Fabiano B *et al*. Occupational injuries in Italy: risk factors and long term trend (1951–98), **330**
Faria MAM, Almeida JWR. Lung cancer mortality in an urban and industrialised area of Brazil: 1980–93: *correspondence*, **136**
Fawell J *see* Keegan T *et al*
Ferrie JE *see* Kivimäki M *et al*
see Kivimäki M *et al*
Feyer A-M *see* Horsburgh S *et al*
Firth-Cozens J *see* Kivimäki M *et al*
Floderus B *see* Nylén L *et al*
see Voss M *et al*
Florvaag E *see* Hollund BE *et al*
Forastiere F *see* Agabiti N *et al*
Foulds IS *see* Chaiear N *et al*
Fradier-Dusch M *see* Nadif R *et al*
Frech W *see* Iregren A *et al*
Freedman M, Burns CJ. Amyotrophic lateral sclerosis and occupational exposure to 2,4-dichlorophenoxyacetic acid: *letter*, **609**
Frumkin H *see* Williams PL *et al*
Fu C *see* Chen W *et al*

Gamberale D *see* Crebelli R *et al*
Gao P *see* Chen W *et al*
Garcia-Gómez M *see* Boffetta P *et al*
Gardiner K *et al*. Respiratory health effects from exposure to carbon black: results of the phase 2 and 3 cross sectional studies in the European carbon black manufacturing industry, **496**
Gawkrodger DJ. Patch testing in occupational dermatology, **823**
Geier J *see* Uter W *et al*
Gefeller O *see* Uter W *et al*
Ghittori S *see* Crebelli R *et al*
Gilbert S *see* Rice FL *et al*
Gillham R *see* Pilkington A *et al*
Glas C *et al*. Hepatitis A in workers exposed to sewage: a systematic review: *review*, **762**
Goldbohm RA *see* Zeegers MPA *et al*
Gomes J *et al*. Dust exposure and impairment of lung function at a small iron foundry in a rapidly developing country, **656**
Goodfellow FJL *et al*. Public health response to an incident of secondary chemical contamination at a beach in the United Kingdom, **232**
Gorman DF *see* Whyte P *et al*
Granath F *et al*. Cancer risk from exposure to occupational acrylamide: *letter*, **608**
Grandi C *see* Iavicoli S *et al*
Griffin MJ *see* Bovenzi M *et al*
see Noël B *et al*
Grimalt J *see* Sala M *et al*
Groat S *see* Cowie HA *et al*
Gustafsson K *see* Iregren A *et al*
Gyntelberg F *see* Suadicani P *et al*

Hagberg S *see* Andersson E *et al*
Hageman G *see* van der Hoek JAF *et al*
Hagman M *see* Iregren A *et al*
Hagmar L *see* Björk J *et al*
Häkkänen M *et al*. Job experience, work load, and risk of musculoskeletal disorders, **129**
Hallermann J *see* Triebig G and Hallermann J
Hanaoka T *see* Zhang J *et al*
Hannerz H, Tüchsen F. Hospital admissions among male drivers in Denmark, **253**
Hansen S *see* Pilkington A *et al*
Hara K *see* Zhang J *et al*
Harrington JM *see* Sorahan T *et al*
Harrington JM *et al*. Leukaemia mortality in relation to magnetic field exposure: findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **307**
Harrington JM. Health effects of shift work and extended hours of work, **68**
Harrington M *see* Gardiner K *et al*
Harrison JC *see* Chen W *et al*
Harrison P. Air pollution research databases: *correspondence*, **64**
Harrison RM *see* Anderson HR *et al*
Hayabuchi H *see* Yoshimura T *et al*
Heederik D *see* Eduard W *et al*
Hein HO *see* Suadicani P *et al*
Heiskel H *see* Seidler A *et al*
Hellier G *see* Nadif R *et al*
Hemingway H *see* Kivimäki M *et al*
Henkel N *see* Seidler A *et al*
Hernandez JC *see* Schulz MR *et al*
Herrero C *see* Sala M *et al*
Hertz-Picciotto I *see* Schulz MR *et al*
Hillerdal G *see* Seldén AI *et al*
Hodgson J *see* Spence A *et al*
Hollund BE *et al*. Prevalence of airway symptoms among hairdressers in Bergen, Norway, **780**
Hong C-S *see* Park JW *et al*
Hoogendoorn WE *see* Ariëns GAM *et al*
Horsburgh S *et al*. Fatal work related injuries in agricultural production and services to agriculture sectors of New Zealand, 1985–94, **489**
Hotz P *see* Glas C *et al*
Howarth S *see* Dennekamp M *et al*
see Dick CAJ *et al*
Hudnell K *see* Engel LS *et al*
Hulshof C *see* Verbeek J *et al*
Hurley JF *see* Buchanan D *et al*
see Cowie HA *et al*
see Pilkington A *et al*
Hwang B-F *see* Jaakkola JJK *et al*

Iavicoli S *et al*. Research priorities in occupational health in Italy, **325**
Ibrahim SA *see* Cole DC *et al*
Ichiba M *see* Zhang J *et al*
Igarashi M *see* Tsutsumi A *et al*

Ihrig A *et al.* Evaluation of a modified German version of the Q16 questionnaire for neurotoxic symptoms in workers exposed to solvents, **19**
Ilmarinen JE. Aging workers, **546**
Imbriani M *see* Crebelli R *et al*
Iregren A *et al.* Effects on the nervous system in different groups of workers exposed to aluminium, **453**
Ismail K. A review of the evidence for a "Gulf war syndrome", **754**
New challenges facing ill health in Gulf war veterans: *editorial*, **289**
Iversen A *see* Goodfellow FJL *et al*
Iversen M *see* Radon K *et al*

Jaakkola JJK *et al.* Foetal growth and duration of gestation relative to water chlorination, **437**
Jamal GA *see* Pilkington A *et al*
James WH. The bones and hormones of deep water divers and pilots of high performance aircraft: *letter*, **682**
Järup L *see* Rosenlund M *et al*
Jeebhay MF *et al.* Occupational seafood allergy: a review, **553**
Joffe M. Are problems with male reproductive health caused by endocrine disruption?, **281**
Johansson B *see* Björk J *et al*
Johnson DL *see* Phillips ML *et al*
Jones M *see* Chaiear N *et al*
see O'Connor TM *et al*

Kaiser U *see* Seidler A *et al*
Kales SN *et al.* Prospective study of hepatic, renal, and haematological surveillance in hazardous materials firefighters, **87**
Kalman CJ. Guidelines on good publication practice: *correspondence*, **137**
Kaneko S *see* Yoshimura T *et al*
Kang DB *see* Park JW *et al*
Kankaanpää E *see* Verbeek J *et al*
Kant I *see* Zeegers MPA *et al*
Karlsson B *et al.* Is there an association between shift work and having a metabolic syndrome? Results from a population based study of 27 485 people, **747**
Kauffmann F *see* Nadif R *et al*
Kauppinen T *see* Björk J *et al*
Kayaba K *see* Tsutsumi A *et al*
Keegan T *et al.* Use of routinely collected data on trihalomethane in drinking water for epidemiological purposes, **447**
Keifer MC *see* Engel LS *et al*
Ketchum N *see* Steenland K *et al*
Kidd M *see* Pilkington A *et al*
Kidd MW *see* Buchanan D *et al*
Kim CW *see* Park JW *et al*
Kim KS *see* Park JW *et al*
King WD *see* Dodds L and King WD
Kirsch-Volders M *see* Lison D *et al*
Kivimäki M *see* Vahtera J *et al*
Kivimäki M *et al.* Organisational downsizing and musculoskeletal problems in employees: a prospective study, **811**
Sickness absence in hospital physicians: 2 year follow up study on determinants, **361**
Knutsson A *see* Karlsson B *et al*
Ko SH *see* Park JW *et al*
Kobal AB *see* Boffetta P *et al*
Koh THHG *see* Koh TS and Koh THHG
Koh TS, Koh THHG. Bullying in hospitals: *letter*, **610**
Kok FJ *see* van Amelsvoort LGPM *et al*
Kokki E *et al.* Small area estimation of incidence of cancer around a known source of exposure with fine resolution data, **315**
Kollmeier J *see* Bauer TT *et al*
Kongerud J *see* Ulvestad B *et al*
Kraus T *et al.* Exposure assessment in the hard metal manufacturing industry with special regard to tungsten and its compounds, **631**
Kurinczuk JJ, Clarke M. Case-control study of leatherwork and male infertility, **217**

Labrèche F *see* Cherry N *et al*
Langley JD *see* Horsburgh S *et al*
Lee AH *see* Carrivick PJW *et al*
Lee B-K *see* Todd AC *et al*
Lee G-S *see* Todd AC *et al*
Lee HS *et al.* Radiological progression and its predictive risk factors in silicosis, **467**
Lee PN. Relation between exposure to asbestos and smoking jointly and the risk of lung cancer: *review*, **145**
Lehrer SB *see* Jeebhay MF *et al*
Lemke B *see* Bauer TT *et al*
Levy L *see* Duarte-Davidson R *et al*
Lindahl B *see* Karlsson B *et al*
Lindgren T *see* Wieslander G *et al*
Lindsell CJ *see* Bovenzi M *et al*
Lison D *et al.* Update on the genotoxicity and carcinogenicity of cobalt compounds, **619**
Ljunggren KG *see* Iregren A *et al*
Lloyd OL *see* Gomes J *et al*
Longstreth Jr WT *see* Engel LS *et al*
Loomis D, Rushton L. Reporting of occupational and environmental research: use and misuse of statistical and epidemiological methods: *correspondence*, **63**
Lopata AL *see* Jeebhay MF *et al*
Lucas LJ *see* Granath F *et al*
Lund MB *see* Ulvestad B *et al*
Lundell D *see* Egedahl R *et al*
Lundgren EAL *see* Seldén AI *et al*
Lundström R *see* Nilsson T and Lundström R
Luukkonen R *see* Vanhanen M *et al*
see Viikari-Juntura E *et al*
Lygre SH *see* Hollund BE *et al*

Maan AC *see* van Amelsvoort LGPM *et al*
McBride DI, Williams S. Audiometric notch as a sign of noise induced hearing loss, **46**
Macfarlane GJ *see* Cherry N *et al*
see Cherry N *et al*
see Nahit ES *et al*
MacNee W *see* Donaldson K *et al*
Maconochie N *see* Doyle P *et al*
Magnus P *see* Jaakkola JJK *et al*
Manca P *see* Carta P *et al*
Manning CC *see* Williams PL *et al*
Mapp CE. Agents, old and new, causing occupational asthma, **354**
Marinaccio A *see* Iavicoli S *et al*
Marsh GM *see* Granath F *et al*
Martikainen R *see* Häkkänen M *et al*
see Miranda H *et al*
see Viikari-Juntura E *et al*
Martini A *see* Crebelli R *et al*
Masschelein R *see* Viaene MK *et al*
Massin N *see* Cowie HA *et al*
Matsuo Y *see* Nakanishi N *et al*
Mauritzson N *see* Björk J *et al*
Meagher CC *see* Phillips ML *et al*
Mehl R *see* Eduard W *et al*
Melbostad E *see* Eduard W *et al*
Mendoza PJ *see* Kales SN *et al*
Merler E *see* Boffetta P *et al*
Meyer HW *see* Suadicani P *et al*
Michalek J *see* Steenland K *et al*
Miedema MC *see* Ariëns GAM *et al*
Mikoczy Z *see* Björk J *et al*
Miller BG *see* Cowie HA *et al*
Miranda H *et al.* A prospective study of work related factors and physical exercise as predictors of shoulder pain, **528**
Mitelman F *see* Björk J *et al*
Mocci F *et al.* Psychological factors and visual fatigue in working with video display terminals, **267**
Moen BE *see* Hollund BE *et al*
Moon J-D *see* Brodkin CA *et al*
Moshier EL *see* Todd AC *et al*
Muirhead CR. Cancer after nuclear incidents, **482**
Murray VSG *see* Goodfellow FJL *et al*
Mutanen P *see* Viikari-Juntura E *et al*

Nadif R *et al.* Cross sectional and longitudinal study on selenium, glutathione peroxidase, smoking, and occupational exposure in coal miners, **239**
Nagano K *see* Nakanishi N *et al*
Nahit ES *et al.* Short term influence of mechanical factors on regional musculoskeletal pain: a study of new workers from 12 occupational groups, **374**
Nakamura K *see* Nakanishi N *et al*
Nakanishi N *et al.* Hours of work and the risk of developing impaired fasting glucose or type 2 diabetes mellitus in Japanese male office workers, **569**
Nemery B. What happens to the manuscripts that have not been accepted for publication in *Occupational and Environmental Medicine*?, **604**
Ng TP *see* Lee HS *et al*
Nichols L *see* Harrington JM *et al*
see Sorahan T *et al*
Nieuwenhuijsen MJ *see* Keegan T *et al*
Nilsson P-G *see* Björk J *et al*
Nilsson T *see* Andersson E *et al*
Nilsson T, Lundström R. Quantitative thermal perception thresholds relative to exposure to vibration, **472**
Nishina K *see* Nakanishi N *et al*
Niven RM *see* Anees W *et al*
Noël B *et al.* Raynaud's phenomenon in workers exposed to vibration: *letter*, **279**
Norbäck D *see* Wieslander G *et al*
Nordman H *see* Vanhanen M *et al*
Norman NJ *see* Gomes J *et al*
Nowak D *see* Radon K *et al*
Nylén L *see* Iregren A *et al*
Nylén L *et al.* Mortality among women and men relative to unemployment, part time work, overtime work, and extra work: a study based on data from the Swedish twin registry, **52**

O'Connor TM *et al.* Report of occupational asthma due to phytase and β-glucanase, **417**
Ohlson C-G *see* Seldén AI *et al*
Ojajärvi A *see* Seilkop SK *et al*
Omenaas E *see* Hollund BE *et al*
Oryszczyn M-P *see* Nadif R *et al*
Osman J *see* Spence A *et al*
Ounsanya T *et al.* Acute respiratory effects of particles: mass or number?, **154**
Ouki SK *see* Goodfellow FJL *et al*

Pahwa P *see* Gomes J *et al*
Palin LA *et al.* Personal exposure of children to nitrogen dioxide: *letter*, **682**
Palmer KT *see* Noël B *et al*
Palmi S *see* Iavicoli S *et al*
Pan G *see* Zhang J *et al*
Panella M *see* Palin LA *et al*
Pannett HB *see* Noël B *et al*
Park JW *et al.* Role of skin prick test and serological measurement of specific IgE in the diagnosis of occupational asthma resulting from exposure to vinyl sulphone reactive dyes, **411**
Park R *see* Rice FL *et al*
Partanen T *see* Seilkop SK *et al*
Pastorino R *see* Fabiano B *et al*
Pattenden S. Air traffic noise and hypertension in Stockholm County: *editorial commentary*, **761**
Paulsson B *see* Granath F *et al*
Pauwels W *see* Viaene MK *et al*
Pedersen S *see* Radon K *et al*
Pekkanen J *see* Kokki E *et al*
Pentti J *see* Kivimäki M *et al*
see Vahtera J *et al*
Penttinen A *see* Kokki E *et al*
Pershagen G *see* Rosenlund M *et al*
Persson B *see* Andersson E *et al*
Perucci CA *see* Agabiti N *et al*
Pfahlberg A *see* Uter W *et al*
Pham Q-T *see* Nadif R *et al*
Phillips ML *et al.* What is "powder free"? Characterisation of powder aerosol produced during simulated use of powdered and powder free latex gloves, **479**
Phoon WH *see* Lee HS *et al*
Pickering CAC *see* Anees W *et al*
Piekarski C *see* Cowie HA *et al*
Pierce ML *see* Williams PL *et al*
Piirainen H *see* Verbeek J *et al*
Pilkington A *see* Buchanan D *et al*
Pilkington A *et al.* An epidemiological study of the relations between exposure to organophosphate pesticides and indices of chronic peripheral neuropathy and neuropsychological abnormalities in sheep farmers and dippers, **702**
Polyhronopoulos GN *see* Kales SN *et al*
Pompe-Kirn V *see* Boffetta P *et al*
Prescott G *see* Osunsanya T *et al*
Presti EL *see* Agabiti N *et al*
Pritchard CM *see* Nahit ES *et al*
Pukkala E *see* Kokki E *et al*
Punnett L *see* Bildt C *et al*

Radon K *et al.* Exposure assessment and lung function in pig and poultry farmers, **405**
Rafnsson V. Incidence of cancer among bookbinders, printers, photoengravers, and typesetters, **523**
Ranta J *see* Kokki E *et al*
Räsänen K *see* Kivimäki M *et al*
see Verbeek J *et al*
Redlich CA *see* Brodkin CA *et al*
Reid K, Dawson D. Comparing performance on a simulated 12 hour shift rotation in young and older subjects, **58**
Renwick L *see* Donaldson K *et al*
Rice FL *et al.* Crystalline silica exposure and lung cancer mortality in diatomaceous earth industry workers: a quantitative risk assessment, **38**
Riihimäki H *see* Miranda H *et al*
see Viikari-Juntura E *et al*
Robertson AS *see* Anees W *et al*
Robins TG *see* Jeebhay MF *et al*
Roels HA *see* Viaene MK *et al*
Roman E *see* Doyle P *et al*
Rosenlund M *et al.* Increased prevalence of hypertension in a population exposed to aircraft noise, **769**
Rothenbacher D *see* Siebert U *et al*
Roth-Küver B *see* Seidler A *et al*
Rushton L *see* Duarte-Davidson R *et al*
see Loomis D and Rushton L

Sadhra S *see* Chaiear N *et al*
Sadhra S *et al.* Occupational health research priorities in Malaysia: a Delphi study, **426**
Sala M *et al.* Association between serum concentrations of hexachlorobenzene and polychlorobiphenyls with thyroid hormone and liver enzymes in a sample of the general population, **172**
Sällsten G *see* Boffetta P *et al*
Sanders AG *see* Williams PL *et al*
Savitz DA. Occupational exposure to magnetic fields and brain cancer: *editorial*, **617**
Schall LC *see* Granath F *et al*
Schaller KH *see* Kraus T *et al*
Schmidt E-W *see* Bauer TT *et al*
Schnuch A *see* Uter W *et al*
Schouten EG *see* van Amelsvoort LGPM *et al*
Schramel P *see* Kraus T *et al*
Schultze-Werninghaus G *see* Bauer TT *et al*
Schulz MR *et al.* Dose-response relation between acrylamide and pancreatic cancer: *letter*, **609**
Schwartz BS *see* Todd AC *et al*
Scott F *see* Cole DC *et al*
Scott KC *see* Engel LS *et al*
Seaton A *see* Chen R *et al*
see Chen R *et al*
see Dennekamp M *et al*
see Dick CAJ *et al*
see Osunsanya T *et al*
Sebastián MS *et al.* Exposures and cancer incidence near oil fields in the Amazon basin of Ecuador, **517**
Seidler A *et al.* The role of cumulative physical work load in lumbar spine disease: risk factors for lumbar osteochondrosis and spondylosis associated with chronic complaints, **735**
Seilkop SK *et al.* Occupational exposures and pancreatic cancer: a meta-analysis: *correspondence*, **63**
Seixas NS *see* Engel LS *et al*
Selcuk T *see* Dumortier P *et al*
Seldén AI *et al.* Exposure to tremolite asbestos and respiratory health in Swedish dolomite workers, **670**
Semple S *see* Chen R *et al*
see Chen R *et al*
Serra A *see* Mocci F *et al*
Sewell C *see* Buchanan D *et al*
Shannon HS *see* Cole DC *et al*
Sheikh-Ahmed K *see* Sadhra S *et al*
Siebert U *et al.* Demonstration of the healthy worker survivor effect in a cohort of workers in the construction industry, **774**
Silman AJ *see* Cherry NM *et al*
see Nahit ES *et al*
Sjögren B *see* Iregren A *et al*
Skerjanc A. Sickness absence in diabetic employees, **432**
Skrondal A *see* Jaakkola JJK *et al*
Smedley J *see* Cherry N *et al*
see Cherry N *et al*
Smith PG *see* Doyle P *et al*
Smith R *see* Rice FL *et al*
Sorahan T *see* Harrington JM *et al*
Sorahan T *et al.* Occupational exposure to magnetic fields relative to mortality from brain tumours: updated and revised findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **626**
Soutar CA *see* Cowie HA *et al*
see Pilkington A *et al*
Sparks A *see* Goodfellow FJL *et al*
Spence A *et al.* Monitoring occupational diseases: response: *editorial*, **692**
Stayner L *see* Rice FL *et al*
Steenland K *et al.* Dioxin and diabetes mellitus: an analysis of the combined NIOSH and Ranch Hand data, **641**
Steffen R *see* Glas C *et al*
Stephens C *see* Sebastián MS *et al*
Stone V *see* Dick CAJ *et al*
see Donaldson K *et al*
Strömberg U *see* Björk J *et al*
Suadicani P *et al.* Exposure to cold and draught, alcohol consumption, and the NS-phenotype are associated with chronic bronchitis: an epidemiological investigation of 3387 men aged 53–75 years: the Copenhagen Male Study, **160**
Suh JH *see* Kales SN *et al*
Sunyer J *see* Sala M *et al*
Sutinen R *see* Kivimäki M *et al*
Suzuki K *see* Nakanishi N *et al*
Swaen GMH *see* Zeegers MPA *et al*
Swenne CA *see* van Amelsvoort LGPM *et al*
Syddall H *see* Noël B *et al*

Takahashi K *see* Zhang J *et al*
Takala E-P *see* Miranda H *et al*
see Viikari-Juntura E *et al*
Tannahill SN *see* Buchanan D *et al*
Tatara K *see* Nakanishi N *et al*
Tavendale R *see* Chen R *et al*
Taylor S *see* Cherry N *et al*
see Cherry N *et al*
Theobald H *see* Bildt C *et al*

Tiikkainen U *see* Vanhanen M *et al*
Tinnerberg H *see* Björk J *et al*
Todd AC *et al*. Predictors of DMSA chelatable lead, tibial lead, and blood lead in 802 Korean lead workers, **73**
To-Figueras J *see* Sala M *et al*
Toledano MB *see* Keegan T *et al*
Tomei F *see* Crebelli R *et al*
Tomokuni K *see* Zhang J *et al*
Topping M. Occupational exposure limits for chemicals, **138**
Torén K *see* Andersson E *et al*
Torgén M *see* Bildt C *et al*
Törnqvist M *see* Granath F *et al*
Töyry S *see* Kivimäki M *et al*
Triebig G *see* Ihrig A *et al*
Triebig G, Hallermann J. Survey of solvent related chronic encephalopathy as an occupational disease in European countries, **575**
Tsutsumi A *et al*. Association between job strain and prevalence of hypertension: a cross sectional analysis in a Japanese working population with a wide range of occupations: the Jichi Medical School cohort study, **367**
Tsutsumi K *see* Tsutsumi A *et al*
Tuchsen F *see* Hannerz H and Tüchsen F
Tulandi T *see* Cherry N *et al*
Tunstall-Pedoe H *see* Chen R *et al*
Tuomainen A *see* Vanhanen M *et al*
Tuomi T *see* Vanhanen M *et al*
Tupasela O *see* Vanhanen M *et al*

Ulvestad B *et al*. Cumulative exposure to dust causes accelerated decline in lung function in tunnel workers, **663**
Ursini CL *see* Iavicoli S *et al*
Uter W *et al*. Association between occupation and contact allergy to the fragrance mix: a multifactorial analysis of national surveillance data, **392**

Vahtera J *see* Kivimäki M *et al*
see Kivimäki M *et al*
Vahtera J *et al*. The role of extended weekends in sickness absenteeism, **818**
Vainio H *see* Wennborg H *et al*
Vale JA *see* Bradberry SM *et al*
van Amelsvoort LGPM *et al*. Changes in frequency of premature complexes and heart rate variability related to shift work, **678**
van den Brandt PA *see* Zeegers MPA *et al*
van der Hoek JAF *et al*. Routine diagnostic procedures for chronic encephalopathy induced by solvents: survey of experts, **382**
van der Laan G *see* van der Hoek JAF *et al*
van der Wal G *see* Ariëns GAM *et al*
van Dijk F *see* Verbeek J *et al*
Vanhanen M *et al*. Sensitisation to enzymes in the animal feed industry, **119**
van Mechelen W *see* Ariëns GAM *et al*
van Tongeren M *see* Gardiner K *et al*
see Harrington JM *et al*
see Sorahan T *et al*
van Wijngaarden E *see* Schulz MR *et al*
Verbeek J *et al*. Consumer satisfaction with occupational health services: should it be measured?, **272**
Verberk MM *see* van der Hoek JAF *et al*
Verougstraete V *see* Lison D *et al*
Veulemans H *see* Viaene MK *et al*
Viaene MK *et al*. Neurobehavioural changes and persistence of complaints in workers exposed to styrene in a polyester boat building plant: influence of exposure characteristics and microsomal epoxide hydrolase phenotype, **103**
Viikari-Juntura E *see* Häkkänen M *et al*
see Miranda H *et al*
Viikari-Juntura E *et al*. Longitudinal study on work related and individual risk factors affecting radiating neck pain, **345**
Vonesch N *see* Iavicoli S *et al*
Voss M *see* Nylén L *et al*
Voss M *et al*. Physical, psychosocial, and organisational factors relative to sickness absence: a study based on Sweden Post, **178**
Vyas A *see* Anees W *et al*

Waclawski ER. Glutaraldehyde induced asthma in endoscopy nursing staff: *letter*, **544**
Walker LG *see* Chen R *et al*
Wallace WE *see* Chen W *et al*
Walters S *see* Anderson HR *et al*
Weber A *see* Bauer TT *et al*
see Kraus T *et al*
Weber C *see* Radon K *et al*
Welinder H *see* Björk J *et al*
Wennberg A *see* Iregren A *et al*
Wennborg H *et al*. Solvent use and time to pregnancy among female personnel in biomedical laboratories in Sweden, **225**
Whitaker H *see* Keegan T *et al*
Whitaker SC. The management of sickness absence, **420**
Whyte P *et al*. Positive reform of tuna farm diving in South Australia in response to government intervention, **124**
Wieslander G *et al*. Experimental exposure to propylene glycol mist in aviation emergency training: acute ocular and respiratory effects, **649**
Wik N-G *see* Seldén AI *et al*
Wikman A *see* Bildt C *et al*
Wild P *see* Cowie HA *et al*
Wilkinson M *see* Keegan T *et al*
Williams NR *see* Bradberry SM *et al*
Williams PL *et al*. Reconstruction of occupational mercury exposures at a chloralkali plant, **81**
Williams S *see* McBride DI and Williams S
Willingstorfer WJ *see* Seidler A *et al*
Willson RA *see* Brodkin CA *et al*
Wingren G *see* Andersson E *et al*
Won JU *see* Park JW *et al*

Yamano Y *see* Zhang J *et al*
Yang JY *see* Park JW *et al*
Yau KKW *see* Carrivick PJW *et al*
Yoshida H *see* Nakanishi N *et al*
Yoshimura T *et al*. Sex ratio in offspring of those affected by dioxin and dioxin-like compounds: the Yusho, Seveso, and Yucheng incidents, **540**
Youk AO *see* Granath F *et al*

Zaridze D *see* Boffetta P *et al*
Zeegers MPA *et al*. Occupational risk factors for male bladder cancer: results from a population based case cohort study in the Netherlands, **590**
Zhang J *et al*. Urinary 1-hydroxypyrene in coke oven workers relative to exposure, alcohol consumption, and metabolic enzymes, **716**
Zhang S *see* Zhang J *et al*
Zhuang Z *see* Chen W *et al*
Zijno A *see* Crebelli R *et al*
Zöbelein P *see* Kraus T *et al*

VOLUME 58 • SUBJECT INDEX

12 hour shifts, Comparing performance on a simulated 12 hour shift rotation in young and older subjects, **58**

accidental exposure, Short term respiratory effects of acute exposure to chlorine due to a swimming pool accident, **399**
acoustic rhinometry, Experimental exposure to propylene glycol mist in aviation emergency training: acute ocular and respiratory effects, **649**
acrylamide, Cancer risk from exposure to occupational acrylamide: *letter*, **608**
Dose-response relation between acrylamide and pancreatic cancer: *letter*, **609**
acrylates, Agents, old and new causing occupational asthma, **354**
aerosol, What is "powder free"? Characterisation of powder aerosol produced during simulated use of powdered and powder free latex gloves, **479**
age, Comparing performance on a simulated 12 hour shift rotation in young and older subjects, **58**
aging work force, Aging workers, **546**
agriculture, Fatal work related injuries in agricultural production and services to agriculture sectors of New Zealand, 1985–94, **489**
air pollution, Acute respiratory effects of particles: mass or number?, **154**
Air pollution research databases: *correspondence*, **64**
air pollution particulates, Particulate matter and daily mortality and hospital admissions in the west midlands conurbation of the United Kingdom: associations with fine and coarse particles, black smoke and sulphate, **504**
air traffic noise, Air traffic noise and hypertension in Stockholm County: *editorial commentary*, **761**
alcohol, Exposure to cold and draught, alcohol consumption, and the NS-phenotype are associated with chronic bronchitis: an epidemiological investigation of 3387 men aged 53–75 years: the Copenhagen Male Study, **160**
alcohol drinking, Urinary 1-hydroxypyrene in coke oven workers relative to exposure, alcohol consumption, and metabolic enzymes, **716**
allergen, Patch testing in occupational dermatology, **823**
allergy, Prevalence of airway symptoms among hairdressers in Bergen, Norway, **780**
Sensitisation to natural rubber latex: an epidemiological study of workers exposed during tapping and glove manufacture in Thailand, **386**
aluminium, Effects on the nervous system in different groups of workers exposed to aluminium, **453**
Amazon, Exposures and cancer incidence near oil fields in the Amazon basin of Ecuador, **517**
aminotransferases, Serum hepatic biochemical activity in two populations of workers exposed to styrene, **95**
amyotrophic lateral sclerosis, Amyotrophic lateral sclerosis and occupational exposure to 2,4-dichlorophenoxyacetic acid: *letter*, **609**
animal feed, Sensitisation to enzymes in the animal feed industry, **119**
anti-androgen, Are problems with male reproductive health caused by endocrine disruption?, **281**
antioxidant enzymes, Cross sectional and longitudinal study on selenium, glutathione peroxidase, smoking, and occupational exposure in coal miners, **239**
aquaculture, Positive reform of tuna farm diving in South Australia in response to government intervention, **124**
arcraft noise, Increased prevalence of hypertension in a population exposed to aircraft noise, **769**
arrhythmia, Changes in frequency of premature complexes and heart rate variability related to shift work, **678**
asbestos, Relation between exposure to asbestos and smoking jointly and the risk of lung cancer: *review*, **145**
asbestos body, Erionite bodies and fibres in bronchoalveolar lavage fluid (BALF) of residents from Tuzköy, Cappadocia, Turkey, **261**
asbestos tremolite, Exposure to tremolite asbestos and respiratory health in Swedish dolomite workers, **670**
assembly work, Job experience, work load, and risk of musculoskeletal disorders, **129**
asthenopia, Psychological factors and visual fatigue in working with video display terminals, **267**
asthma, Glutaraldehyde induced asthma in endoscopy nursing staff: *letter*, **544**
Personal exposure of children to nitrogen dioxide: *letter*, **682**
Prevalence of airway symptoms among hairdressers in Bergen, Norway, **780**
aviation medicine, Experimental exposure to propylene glycol mist in aviation emergency training: acute ocular and respiratory effects, **649**
back pain, Work correlates of back problems and activity restriction due to musculoskeletal disorders in the Canadian national population health survey (NPHS) 1994–5 data, **728**
benzene, Benzene in the environment: an assessment of the potential risks to the health of the population: *review*, **2**
benzofurans, Sex ratio in offspring of those affected by dioxin and dioxin-like compounds: the Yusho, Seveso, and Yucheng incidents, **540**
bibliometry, What happens to the manuscripts that have not been accepted for publication in *Occupational and Environmental Medicine*?, **604**
bilirubin, Serum hepatic biochemical activity in two populations of workers exposed to styrene, **95**
bioaerosols, Short term exposure to airborne microbial agents during farm work: exposure-response relations with eye and respiratory symptoms, **113**
biological monitoring, Exposure assessment in the hard metal manufacturing industry with special regard to tungsten and its compounds, **631**
biomonitoring, Exposure to benzene in urban workers: environmental and biological monitoring of traffic police in Rome, **165**
birth defects, Relation between trihalomethane compounds and birth defects, **443**
birth outcomes, Use of routinely collected data on trihalomethane in drinking water for epidemiological purposes, **447**
bladder cancer, Occupational risk factors for male bladder cancer: results from a population based case cohort study in the Netherlands, **590**
bone lead, Predictors of DMSA chelatable lead, tibial lead, and blood lead in 802 Korean lead workers, **73**
bones, The bones and hormones of deep water divers and pilots of high performance aircraft: *letter*, **682**
brain cancer, Occupational exposure to magnetic fields and brain cancer: *editorial*, **617**
brain neoplasm, A case-referent study of cancer mortality among sulfate mill workers in Sweden, **321**
brain tumours, Occupational exposure to magnetic fields relative to mortality from brain tumours: updated and revised findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **626**
Brazil, Lung cancer mortality in an urban and industrialised area of Brazil: 1980–93: *correspondence*, **136**
bronchial obstruction, Functional variables associated with the clinical grade of dyspnoea in coal miners with pneumoconiosis and mild bronchial obstruction, **794**
bullying, Bullying in hospitals: *letter*, **610**

Canada, Work correlates of back problems and activity restriction due to musculoskeletal disorders in the Canadian national population health survey (NPHS) 1994–5 data, **728**
cancer, Cancer after nuclear incidents, **482**
Cancer risk from exposure to occupational acrylamide: *letter*, **608**
Exposures and cancer incidence near oil fields in the Amazon basin of Ecuador, **517**
carbon black, Respiratory health effects from exposure to carbon black: results of the phase 2 and 3 cross sectional studies in the European carbon black manufacturing industry, **496**
cardiovascular diseases, Mortality from cardiovascular diseases and exposure to inorganic mercury, **461**
case-control study, Are occupational, hobby, or lifestyle exposures associated with Philadelphia chromosome positive chronic myeloid leukaemia?, **722**
The role of cumulative physical work load in lumbar spine disease: risk factors for lumbar osteochondrosis and spondylosis associated with chronic complaints, **735**
causes of cancer, Defamatory article by Martin Walker: *correspondence*, **136**
ceramic fibres, An epidemiological study of the respiratory health of workers in the European refractory ceramic fibre industry, **800**
chemical, Public health response to an incident of secondary chemical contamination at a beach in the United Kingdom, **232**
chemicals, Occupational exposure limits for chemicals, **138**
children, Personal exposure of children to nitrogen dioxide: *letter*, **682**
chloralkali, Reconstruction of occupational mercury exposures at a chloralkali plant, **81**
chlorination byproducts, Use of routinely collected data on trihalomethane in drinking water for epidemiological purposes, **447**
chlorine gas, Short term respiratory effects of acute exposure to chlorine due to a swimming pool accident, **399**
chronic bronchitis, Exposure to cold and draught, alcohol consumption, and the NS-phenotype are associated with chronic bronchitis: an epidemiological investigation of 3387 men aged 53–75 years: the Copenhagen Male Study, **160**
chronic obstructive lung disease, Acute respiratory effects of particles: mass or number?, **154**

chronic toxic encephalopathy, Routine diagnostic procedures for chronic encephalopathy induced by solvents: survey of experts, **382**
circadian rhythms, Comparing performance on a simulated 12 hour shift rotation in young and older subjects, **58**
classification, Routine diagnostic procedures for chronic encephalopathy induced by solvents: survey of experts, **382**
clusters, Health and exposures of United Kingdom Gulf war veterans. Part I: The pattern and extent of ill health, **291**
coal miners, Cross sectional and longitudinal study on selenium, glutathione peroxidase, smoking, and occupational exposure in coal miners, **239**
coal workers' pneumoconiosis, Functional variables associated with the clinical grade of dyspnoea in coal miners with pneumoconiosis and mild bronchial obstruction, **794**
cobalt, Exposure assessment in the hard metal manufacturing industry with special regard to tungsten and its compounds, **631**
Update on the genotoxicity and carcinogenicity of cobalt compounds, **619**
cohort mortality study, Leukaemia mortality in relation to magnetic field exposure: findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **307**
Occupational exposure to magnetic fields relative to mortality from brain tumours: updated and revised findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **626**
cold, Exposure to cold and draught, alcohol consumption, and the NS-phenotype are associated with chronic bronchitis: an epidemiological investigation of 3387 men aged 53–75 years: the Copenhagen Male Study, **160**
construction workers, Demonstration of the healthy worker survivor effect in a cohort of workers in the construction industry, **774**
consumer satisfaction, Consumer satisfaction with occupational health services: should it be measured?, **272**
contact allergy, Association between occupation and contact allergy to the fragrance mix: a multifactorial analysis of national surveillance data, **392**
continuing professional development, Continuing Professional Development in *Occupational and Environmental Medicine*: *editorial*, **1**
cooking fuels, Ultrafine particles and nitrogen oxides generated by gas and electric cooking, **511**
coronary heart disease, Is there an association between shift work and having a metabolic syndrome? Results from a population based study of 27 485 people, **747**
cristobalite, Crystalline silica exposure and lung cancer mortality in diatomaceous earth industry workers: a quantitative risk assessment, **38**
cross sectional survey, Work correlates of back problems and activity restriction due to musculoskeletal disorders in the Canadian national population health survey (NPHS) 1994–5 data, **728**
cryptorchidism, Are problems with male reproductive health caused by endocrine disruption?, **281**
crystalline silica, Crystalline silica exposure and lung cancer mortality in diatomaceous earth industry workers: a quantitative risk assessment, **38**
Mortality from lung cancer among silicotic patients in Sardinia: an update study with 10 more years of follow up, **786**
cyanides, Public health response to an incident of secondary chemical contamination at a beach in the United Kingdom, **232**

database, Air pollution research databases: *correspondence*, **64**
Declaration of Helsinki, Ethics in occupational health research, **685**
decompression illness, Positive reform of tuna farm diving in South Australia in response to government intervention, **124**
developing countries, Getting *Occupational and Environmental Medicine* to developing countries: *editorial*, **425**
diabetes, Dioxin and diabetes mellitus: an analysis of the combined NIOSH and Ranch Hand data, **641**
diabetes mellitus, Sickness absence in diabetic employees, **432**
diagnostic criteria, Routine diagnostic procedures for chronic encephalopathy induced by solvents: survey of experts, **382**
2,4-dichlorophenoxyacetic acid, Amyotrophic lateral sclerosis and occupational exposure to 2,4-dichlorophenoxyacetic acid: *letter*, **609**
Mortality in chemical workers potentially exposed to 2,4-dichlorophenoxyacetic acid (2,4-D) 1945–94: an update, **24**
dimercaptosuccinic acid, Predictors of DMSA chelatable lead, tibial lead, and blood lead in 802 Korean lead workers, **73**
dioxin, Dioxin and diabetes mellitus: an analysis of the combined NIOSH and Ranch Hand data, **641**
disease mapping, Small area estimation of incidence of cancer around a known source of exposure with fine resolution data, **315**
DNA breakage, Update on the genotoxicity and carcinogenicity of cobalt compounds, **619**
dockyard painting, Nasal, eye, and skin irritation in dockyard painters, **542**
dockyards, Exposure to organic solvents and personality, **14**
dolomite, Exposure to tremolite asbestos and respiratory health in Swedish dolomite workers, **670**
dose-response relation, Exposure to silica and silicosis among tin miners in China: exposure-response analyses and risk assessment, **31**
downsizing, Organisational downsizing and musculoskeletal problems in employees: a prospective study, **811**
draught, Exposure to cold and draught, alcohol consumption, and the NS-phenotype are associated with chronic bronchitis: an epidemiological investigation of 3387 men aged 53–75 years: the Copenhagen Male Study, **160**
dust exposure, Cross sectional and longitudinal study on selenium, glutathione peroxidase, smoking, and occupational exposure in coal miners, **239**
Dust exposure and impairment of lung function at a small iron foundry in a rapidly developing country, **656**

early retirement, Demonstration of the healthy worker survivor effect in a cohort of workers in the construction industry, **774**
Ecuador, Exposures and cancer incidence near oil fields in the Amazon basin of Ecuador, **517**
education, Positive reform of tuna farm diving in South Australia in response to government intervention, **124**
electricity generation and transmission, Leukaemia mortality in relation to magnetic field exposure: findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **307**
Occupational exposure to magnetic fields relative to mortality from brain tumours: updated and revised findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **626**
employment, Aging workers, **546**
encephalopathy, Survey of solvent related chronic encephalopathy as an occupational disease in European countries, **575**
energy equivalent acceleration magnitude, Response of finger circulation to energy equivalent combinations of magnitude and duration of vibration, **185**
environment, Benzene in the environment: an assessment of the potential risks to the health of the population: *review*, **2**
Lung cancer mortality in an urban and industrialised area of Brazil: 1980–93: *correspondence*, **136**
environmental exposure, Association between serum concentrations of hexachlorobenzene and polychlorobiphenyls with thyroid hormone and liver enzymes in a sample of the general population, **172**
Erionite bodies and fibres in bronchoalveolar lavage fluid (BALF) of residents from Tuzköy, Cappadocia, Turkey, **261**
enzyme allergy, Sensitisation to enzymes in the animal feed industry, **119**
enzymes, Agents, old and new, causing occupationalasthma, **354**
epidemiology, A case-referent study of cancer mortality among sulfate mill workers in Sweden, **321**
Are occupational, hobby, or lifestyle exposures associated with Philadelphia chromosome positive chronic myeloid leukaemia?, **722**
Mortality experience among employees at a hydrometallurgical nickel refinery and fertiliser complex in Fort Saskatchewan, Alberta (1954–95), **711**
Mortality from cardiovascular diseases and exposure to inorganic mercury, **461**
Occupational risk factors for male bladder cancer: results from a population based case cohort study in the Netherlands, **590**
Short term influence of mechanical factors on regional musculoskeletal pain: a study of new workers from 12 occupational groups, **374**
ergonomics, Consultative team to assess manual handling and reduce the risk of occupational injury, **339**
ethics, Ethics in occupational health research, **685**
European multicentre study, Exposure assessment and lung function in pig and poultry farmers, **405**
exposure, Short term exposure to airborne microbial agents during farm work: exposure-response relations with eye and respiratory symptoms, **113**
Use of routinely collected data on trihalomethane in drinking water for epidemiological purposes, **447**
exposure model, Estimation of cumulative exposure to organophosphate sheep dips in a study of chronic neurological health effects among United Kingdom sheep dippers, **694**
exposure to benzene, Exposure to benzene in urban workers: environmental and biological monitoring of traffic police in Rome, **165**
exposure to silica, Exposure to silica and silicosis among tin miners in China: exposure-response analyses and risk assessment, **31**
exposure-response, Short term exposure to airborne microbial agents during farm work: exposure-response relations with eye and respiratory symptoms, **113**

farmer, Parkinsonism and occupational exposure to pesticides, **582**
fatal injury, Fatal work related injuries in agricultural production and services to agriculture sectors of New Zealand, 1985–94, **489**
ferruginous body, Erionite bodies and fibres in bronchoalveolar lavage fluid (BALF) of residents from Tuzköy, Cappadocia, Turkey, **261**

finger circulation, Response of finger circulation to energy equivalent combinations of magnitude and duration of vibration, **185**
firefighters, Prospective study of hepatic, renal, and haematological surveillance in hazardous materials firefighters, **87**
flour, Agents, old and new, causing occupational asthma, **354**
foetal growth, Foetal growth and duration of gestation relative to water chlorination, **437**
foundry, Dust exposure and impairment of lung function at a small iron foundry in a rapidly developing country, **656**
fragrances, Association between occupation and contact allergy to the fragrance mix: a multifactorial analysis of national surveillance data, **392**
fungal spores, Short term exposure to airborne microbial agents during farm work: exposure-response relations with eye and respiratory symptoms, **113**

genetic marker, Exposure to cold and draught, alcohol consumption, and the NS-phenotype are associated with chronic bronchitis: an epidemiological investigation of 3387 men aged 53–75 years: the Copenhagen Male Study, **160**
genetic polymorphism, Urinary 1-hydroxypyrene in coke oven workers relative to exposure, alcohol consumption, and metabolic enzymes, **716**
gestational age, Foetal growth and duration of gestation relative to water chlorination, **437**
glove factory, Sensitisation to natural rubber latex: an epidemiological study of workers exposed during tapping and glove manufacture in Thailand, **386**
glove powder, What is "powder free"? Characterisation of powder aerosol produced during simulated use of powdered and powder free latex gloves, **479**
β-glucanase Report of occupational asthma due to phytase and β-glucanase, **417**
glutaraldehyde, Agents, old and new, causing occupational asthma, **354**
granite quarry, Radiological progression and its predictive risk factors in silicosis, **467**
guidelines, Guidelines on good publication practice: *correspondence*, **137**
Gulf war, Health and exposures of United Kingdom Gulf war veterans. Part I: The pattern and extent of ill health, **291**
Health and exposures of United Kingdom Gulf war veterans. Part II: The relation of health to exposure, **299**
New challenges facing ill health in Gulf war veterans: *editorial*, **289**
Gulf war syndrome, A review of the evidence for a "Gulf war syndrome", **754**

hairdresser, Prevalence of airway symptoms among hairdressers in Bergen, Norway, **780**
hazardous materials, Prospective study of hepatic, renal, and haematological surveillance in hazardous materials firefighters, **87**
health care personnel, Sickness absence in hospital physicians: 2 year follow up study on determinants, **361**
health effect, Nasal, eye, and skin irritation in dockyard painters, **542**
healthcare, Sensitisation to natural rubber latex: an epidemiological study of workers exposed during tapping and glove manufacture in Thailand, **386**
healthy worker survivor effect, Demonstration of the healthy worker survivor effect in a cohort of workers in the construction industry, **774**
hearing loss, Audiometric notch as a sign of noise induced hearing loss, **46**
heart rate, Changes in frequency of premature complexes and heart rate variability related to shift work, **678**
heat pain, Quantitative thermal perception thresholds relative to exposure to vibration, **472**
heavy construction, Cumulative exposure to dust causes accelerated decline in lung function in tunnel workers, **663**
hepatitis A, Hepatitis A in workers exposed to sewage: a systematic review: *review*, **762**
hepatotoxicity, Serum hepatic biochemical activity in two populations of workers exposed to styrene, **95**
hexachlorobenzene, Association between serum concentrations of hexachlorobenzene and polychlorobiphenyls with thyroid hormone and liver enzymes in a sample of the general population, **172**
hormones, The bones and hormones of deep water divers and pilots of high performance aircraft: *letter*, **682**
hospital admissions, Hospital admissions among male drivers in Denmark, **253**
Particulate matter and daily mortality and hospital admissions in the west midlands conurbation of the United Kingdom: associations with fine and coarse particles, black smoke and sulphate, **504**
hospitals, Bullying in hospitals: *letter*, **610**
hydrocarbons, Public health response to an incident of secondary chemical contamination at a beach in the United Kingdom, **232**
1-hydroxypyrene, Urinary 1-hydroxypyrene in coke oven workers relative to exposure, alcohol consumption, and metabolic enzymes, **716**
hypertension, Air traffic noise and hypertension in Stockholm County: *editorial commentary*, **761**
Association between job strain and prevalence of hypertension: a cross sectional analysis in a Japanese working population with a wide range of occupations: the Jichi Medical School cohort study, **367**
Increased prevalence of hypertension in a population exposed to aircraft noise, **769**
hypospadias, Are problems with male reproductive health caused by endocrine disruption?, **281**

impact factor, What happens to the manuscripts that have not been accepted for publication in *Occupational and Environmental Medicine*?, **604**
impaired fasting glucose, Hours of work and the risk of developing impaired fasting glucose or type 2 diabetes mellitus in Japanese male office workers, **569**
incidence of sickness, Physical, psychosocial, and organisational factors relative to sickness absence: a study based on Sweden Post, **178**
indoor air pollution, Stimulation of IL-8 release from epithelial cells by gas cooker PM_{10}: a pilot study, **208**
infertility, Occupational exposure to solvents and male infertility, **635**
Primary infertility in nuclear industry employees: report from the nuclear industry family study, **535**
inhibition of DNA repair, Update on the genotoxicity and carcinogenicity of cobalt compounds, **619**
injury, Consultative team to assess manual handling and reduce the risk of occupational injury, **339**
inoculations, Health and exposures of United Kingdom Gulf war veterans. Part II: The relation of health to exposure, **299**
inorganic mercury, Mortality from cardiovascular diseases and exposure to inorganic mercury, **461**
interleukin-8, Stimulation of IL-8 release from epithelial cells by gas cooker PM_{10}: a pilot study, **208**
interrelations, Interrelations of risk factors and low back pain in scaffolders, **597**
irritant, Patch testing in occupational dermatology, **823**
irritant symptoms, Nasal, eye, and skin irritation in dockyard painters, **542**
isocyanates, Agents, old and new, causing occupational asthma, **354**

job exposure matrix, Reconstruction of occupational mercury exposures at a chloralkali plant, **81**
job strain, Work correlates of back problems and activity restriction due to musculoskeletal disorders in the Canadian national population health survey (NPHS) 1994–5 data, **728**
journalology, What happens to the manuscripts that have not been accepted for publication in *Occupational and Environmental Medicine*?, **604**

laboratory work, Solvent use and time to pregnancy among female personnel in biomedical laboratories in Sweden, **225**
labour injuries, Occupational injuries in Italy: risk factors and long term trend (1951–98), **330**
latex, Agents, old and new, causing occupational asthma, **354**
Sensitisation to natural rubber latex: an epidemiological study of workers exposed during tapping and glove manufacture in Thailand, **386**
latex gloves, What is "powder free"? Characterisation of powder aerosol produced during simulated use of powdered and powder free latex gloves, **479**
leatherwork, Case-control study of leatherwork and male infertility, **217**
leukaemia, Leukaemia mortality in relation to magnetic field exposure: findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **307**
liver enzymes, Association between serum concentrations of hexachlorobenzene and polychlorobiphenyls with thyroid hormone and liver enzymes in a sample of the general population, **172**
long overtime, Hours of work and the risk of developing impaired fasting glucose or type 2 diabetes mellitus in Japanese male office workers, **569**
longitudinal cohort study, Are neck flexion, neck rotation, and sitting at work risk factors for neck pain? Results of a prospective cohort study, **200**
low back pain, Interrelations of risk factors and low back pain in scaffolders, **597**
lumbar osteochondrosis, The role of cumulative physical work load in lumbar spine disease: risk factors for lumbar osteochondrosis and spondylosis associated with chronic complaints, **735**
lumbar spondylosis, The role of cumulative physical work load in lumbar spine disease: risk factors for lumbar osteochondrosis and spondylosis associated with chronic complaints, **735**
lung, Ultrafine particles, **211**
lung cancer, Crystalline silica exposure and lung cancer mortality in diatomaceous earth industry workers: a quantitative risk assessment, **38**
Lung cancer mortality in an urban and industrialised area of Brazil: 1980–93: *correspondence*, **136**
lung cancer mortality, Mortality from lung cancer among silicotic patients in Sardinia: an update study with 10 more years of follow up, **786**

lung function, Cumulative exposure to dust causes accelerated decline in lung function in tunnel workers, **663**
Dust exposure and impairment of lung function at a small iron foundry in a rapidly developing country, **656**
Environmental tobacco smoke and lung function in employees who never smoked: the Scottish MONICA study, **563**
Exposure to tremolite asbestos and respiratory health in Swedish dolomite workers, **670**
Short term respiratory effects of acute exposure to chlorine due to a swimming pool accident, **399**
lung neoplasms, Relation between exposure to asbestos and smoking jointly and the risk of lung cancer: *review*, **145**

magnetic fields, Leukaemia mortality in relation to magnetic field exposure: findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **307**
Occupational exposure to magnetic fields and brain cancer: *editorial*, **617**
Occupational exposure to magnetic fields relative to mortality from brain tumours: updated and revised findings from a study of United Kingdom electricity generation and transmission workers, 1973–97, **626**
Malaysia, Occupational health research priorities in Malaysia: a Delphi study, **426**
male infertility, Case-control study of leatherwork and male infertility, **217**
male manual workers, Occupational exposure to solvents and male infertility, **635**
management, The role of extended weekends in sickness absenteeism, **818**
manual handling, Short term influence of mechanical factors on regional musculoskeletal pain: a study of new workers from 12 occupational groups, **374**
medical research, Ethics in occupational health research, **685**
medical surveillance, Prospective study of hepatic, renal, and haematological surveillance in hazardous materials firefighters, **87**
mental functional capacity, Aging workers, **546**
mental stress, A prospective study of work related factors and physical exercise as predictors of shoulder pain, **528**
Longitudinal study on work related and individual risk factors affecting radiating neck pain, **345**
mercury, Reconstruction of occupational mercury exposures at a chloralkali plant, **81**
mesothelioma, Erionite bodies and fibres in bronchoalveolar lavage fluid (BALF) of residents from Tuzköy, Cappadocia, Turkey, **261**
meta-analysis, Occupational exposures and pancreatic cancer: a meta-analysis: *correspondence*, **63**
metabolic variables, Is there an association between shift work and having a metabolic syndrome? Results from a population based study of 27 485 people, **747**
methaemoglobinaemia, Occupational methaemoglobinaemia, **611**
methodological study, Effects of drop out in a longitudinal study of musculoskeletal disorders, **194**
methods, Reporting of occupational and environmental research: use and misuse of statistical and epidemiological methods: *correspondence*, **63**
methylene blue, Occupational methaemoglobinaemia, **611**
micro-organism, Exposure assessment and lung function in pig and poultry farmers, **405**
microsomal epoxide hydrolase, Neurobehavioural changes and persistence of complaints in workers exposed to styrene in a polyester boat building plant: influence of exposure characteristics and microsomal epoxide hydrolase phenotype, **103**
mining, Mortality from cardiovascular diseases and exposure to inorganic mercury, **461**
MNS, Exposure to cold and draught, alcohol consumption, and the NS-phenotype are associated with chronic bronchitis: an epidemiological investigation of 3387 men aged 53–75 years: the Copenhagen Male Study, **160**
mortality, Mortality among women and men relative to unemployment, part time work, overtime work, and extra work: a study based on data from the Swedish twin registry, **52**
Mortality experience among employees at a hydrometallurgical nickel refinery and fertiliser complex in Fort Saskatchewan, Alberta (1954–95), **711**
Particulate matter and daily mortality and hospital admissions in the west midlands conurbation of the United Kingdom: associations with fine and coarse particles, black smoke and sulphate, **504**
multisystem symptoms, A review of the evidence for a "Gulf war syndrome", **754**
musculoskeletal disorders, Organisational downsizing and musculoskeletal problems in employees: a prospective study, **811**

natural rubber, Sensitisation to natural rubber latex: an epidemiological study of workers exposed during tapping and glove manufacture in Thailand, **386**
neck disorder, Longitudinal study on work related and individual risk factors affecting radiating neck pain, **345**
neck pain, Are neck flexion, neck rotation, and sitting at work risk factors for neck pain? Results of a prospective cohort study, **200**
nervous system, Effects on the nervous system in different groups of workers exposed to aluminium, **453**
neurobehavioural performance, Neurobehavioural changes and persistence of complaints in workers exposed to styrene in a polyester boat building plant: influence of exposure characteristics and microsomal epoxide hydrolase phenotype, **103**
neuropathy, An epidemiological study of the relations between exposure to organophosphate pesticides and indices of chronic peripheral neuropathy and neuropsychological abnormalities in sheep farmers and dippers, **702**
neuroticism, Exposure to organic solvents and personality, **14**
neurotoxicity, Evaluation of a modified German version of the Q16 questionnaire for neurotoxic symptoms in workers exposed to solvents, **19**
Neurobehavioural changes and persistence of complaints in workers exposed to styrene in a polyester boat building plant: influence of exposure characteristics and microsomal epoxide hydrolase phenotype, **103**
new workers, Job experience, work load, and risk of musculoskeletal disorders, **129**
nickel workers, Mortality experience among employees at a hydrometallurgical nickel refinery and fertiliser complex in Fort Saskatchewan, Alberta (1954–95), **711**
night work, Comparing performance on a simulated 12 hour shift rotation in young and older subjects, **58**
nitrogen dioxide, Personal exposure of children to nitrogen dioxide: *letter*, **682**
nitrogen oxides, Ultrafine particles and nitrogen oxides generated by gas and electric cooking, **511**
noise, Audiometric notch as a sign of noise induced hearing loss, **46**
noise induced, Audiometric notch as a sign of noise induced hearing loss, **46**
non-Hodgkin's lymphoma, Incidence of cancer among bookbinders, printers, photoengravers, and typesetters, **523**
Mortality in chemical workers potentially exposed to 2,4-dichlorophenoxyacetic acid (2,4-D) 1945–94: an update, **24**
non-response, Effects of drop out in a longitudinal study of musculoskeletal disorders, **194**
nuclear accidents, Cancer after nuclear incidents, **482**
nuclear industry, Primary infertility in nuclear industry employees: report from the nuclear industry family study, **535**
nurses, Glutaraldehyde induced asthma in endoscopy nursing staff: *letter*, **544**

occupation, Mortality in chemical workers potentially exposed to 2,4-dichlorophenoxyacetic acid (2,4-D) 1945–94: an update, **24**
Occupational risk factors for male bladder cancer: results from a population based case cohort study in the Netherlands, **590**
occupational, Audiometric notch as a sign of noise induced hearing loss, **46**
occupational and environmental medicine, Continuing Professional Development in *Occupational and Environmental Medicine*: *editorial*, **1**
occupational asthma, Agents, old and new, causing occupational asthma, **354**
Occupational seafood allergy: a review, **553**
Report of occupational asthma due to phytase and β-glucanase, **417**
Role of skin prick test and serological measurement of specific IgE in the diagnosis of occupational asthma resulting from exposure to vinyl sulphone reactive dyes, **411**
occupational dermatitis, Patch testing in occupational dermatology, **823**
occupational disease, Survey of solvent related chronic encephalopathy as an occupational disease in European countries, **575**
occupational exposure, Amyotrophic lateral sclerosis and occupational exposure to 2,4-dichlorophenoxyacetic acid: *letter*, **609**
Cancer risk from exposure to occupational acrylamide: *letter*, **608**
Exposure to cold and draught, alcohol consumption, and the NS-phenotype are associated with chronic bronchitis: an epidemiological investigation of 3387 men aged 53–75 years: the Copenhagen Male Study, **160**
Occupational exposures and pancreatic cancer: a meta-analysis: *correspondence*, **63**
Reconstruction of occupational mercury exposures at a chloralkali plant, **81**
Sensitisation to enzymes in the animal feed industry, **119**
occupational exposure limits, Occupational exposure limits for chemicals, **138**
occupational health, Occupational health research priorities in Malaysia: a Delphi study, **426**
Research priorities in occupational health in Italy, **325**
Sickness absence in hospital physicians: 2 year follow up study on determinants, **361**
The role of extended weekends in sickness absenteeism, **818**
occupational health services, Consumer satisfaction with occupational health services: should it be measured?, **272**
occupational illness, Monitoring occupational diseases: response: *editorial*, **692**
Monitoring trends in occupational illness: *editorial*, **691**

occupational mobility, Demonstration of the healthy worker survivor effect in a cohort of workers in the construction industry, 774
occupational risk factors, Association between occupation and contact allergy to the fragrance mix: a multifactorial analysis of national surveillance data, 392
occupational seafood allergy, Occupational seafood allergy: a review, 553
ocupational health, Organisational downsizing and musculoskeletal problems in employees: a prospective study, 811
oestrogen, Are problems with male reproductive health caused by endocrine disruption?, 281
oil, Exposures and cancer incidence near oil fields in the Amazon basin of Ecuador, 517
organic solvents, Exposure to organic solvents and personality, 14
Solvent use and time to pregnancy among female personnel in biomedical laboratories in Sweden, 225
Survey of solvent related chronic encephalopathy as an occupational disease in European countries, 575
organophosphate pesticides, Estimation of cumulative exposure to organophosphate sheep dips in a study of chronic neurological health effects among United Kingdom sheep dippers, 694
organophosphates, Adverse health effects after low level exposure to organophosphates: *editorial*, 689
An epidemiological study of the relations between exposure to organophosphate pesticides and indices of chronic peripheral neuropathy and neuropsychological abnormalities in sheep farmers and dippers, 702
overtime work, Health effects of shift work and extended hours of work, 68
Mortality among women and men relative to unemployment, part time work, overtime work, and extra work: a study based on data from the Swedish twin registry, 52
oxidation, Occupational methaemoglobinaemia, 611

paint exposures, Nasal, eye, and skin irritation in dockyard painters, 542
painter, Exposure to organic solvents and personality, 14
pancreatic cancer, Dose-response relation between acrylamide and pancreatic cancer: *letter*, 609
Occupational exposures and pancreatic cancer: a meta-analysis: *correspondence*, 63
panel study, Effects of drop out in a longitudinal study of musculoskeletal disorders, 194
parkinsonism, Parkinsonism and occupational exposure to pesticides, 582
participatory, Consultative team to assess manual handling and reduce the risk of occupational injury, 339
passive smoking, Environmental tobacco smoke and lung function in employees who never smoked: the Scottish MONICA study, 563
patch testing, Patch testing in occupational dermatology, 823
performance, Comparing performance on a simulated 12 hour shift rotation in young and older subjects, 58
personal protection, Dust exposure and impairment of lung function at a small iron foundry in a rapidly developing country, 656
personality, Exposure to organic solvents and personality, 14
pesticides, Health and exposures of United Kingdom Gulf war veterans. Part II: The relation of health to exposure, 299
Parkinsonism and occupational exposure to pesticides, 582
phenoxy herbicide, Mortality in chemical workers potentially exposed to 2,4-dichlorophenoxyacetic acid (2,4-D) 1945–94: an update, 24
physical risk factors, Are neck flexion, neck rotation, and sitting at work risk factors for neck pain? Results of a prospective cohort study, 200
physical work capacity, Aging workers, 546
physical work load, A prospective study of work related factors and physical exercise as predictors of shoulder pain, 528
Job experience, work load, and risk of musculoskeletal disorders, 129
Longitudinal study on work related and individual risk factors affecting radiating neck pain, 345
The role of cumulative physical work load in lumbar spine disease: risk factors for lumbar osteochondrosis and spondylosis associated with chronic complaints, 735
phytase, Report of occupational asthma due to phytase and β-glucanase, 417
PM_{10}, Stimulation of IL-8 release from epithelial cells by gas cooker PM_{10}: a pilot study, 208
point source, Small area estimation of incidence of cancer around a known source of exposure with fine resolution data, 315
polychlorinated biphenyls, Sex ratio in offspring of those affected by dioxin and dioxin-like compounds: the Yusho, Seveso, and Yucheng incidents, 540
pressure, The bones and hormones of deep water divers and pilots of high performance aircraft: *letter*, 682
printing industry, Incidence of cancer among bookbinders, printers, photoengravers, and typesetters, 523
professional, Positive reform of tuna farm diving in South Australia in response to government intervention, 124
professional drivers, Hospital admissions among male drivers in Denmark, 253
propylene glycol, Experimental exposure to propylene glycol mist in aviation emergency training: acute ocular and respiratory effects, 649
protein contact dermatitis, Occupational seafood allergy: a review, 553
psychological, Association between job strain and prevalence of hypertension: a cross sectional analysis in a Japanese working population with a wide range of occupations: the Jichi Medical School cohort study, 367
psychosocial factors, Sickness absence in hospital physicians: 2 year follow up study on determinants, 361
public health, Public health response to an incident of secondary chemical contamination at a beach in the United Kingdom, 232
publications, Getting *Occupational and Environmental Medicine* to developing countries: *editorial*, 425
publishing, Guidelines on good publication practice: *correspondence*, 137
pulp and paper industry, A case-referent study of cancer mortality among sulfate mill workers in Sweden, 321

quantitative thermal perception, Quantitative thermal perception thresholds relative to exposure to vibration, 472
questionnaire, Short term influence of mechanical factors on regional musculoskeletal pain: a study of new workers from 12 occupational groups, 374
questionnaire Q18, Evaluation of a modified German version of the Q16 questionnaire for neurotoxic symptoms in workers exposed to solvents, 19

radiation, Defamatory article by Martin Walker: *correspondence*, 136
radiation exposure, Cancer after nuclear incidents, 482
radiological progression, Radiological progression and its predictive risk factors in silicosis, 467
Raynaud's phenomenon, Raynaud's phenomenon in workers exposed to vibration: *letter*, 279
reactive dye, Role of skin prick test and serological measurement of specific IgE in the diagnosis of occupational asthma resulting from exposure to vinyl sulphone reactive dyes, 411
reactive oxygen species, Cross sectional and longitudinal study on selenium, glutathione peroxidase, smoking, and occupational exposure in coal miners, 239
regression model, Occupational injuries in Italy: risk factors and long term trend (1951–98), 330
reproductive development, Are problems with male reproductive health caused by endocrine disruption?, 281
research, Air pollution research databases: *correspondence*, 64
Reporting of occupational and environmental research: use and misuse of statistical and epidemiological methods: *correspondence*, 63
research priorities, Occupational health research priorities in Malaysia: a Delphi study, 426
Research priorities in occupational health in Italy, 325
respirable dust, Cumulative exposure to dust causes accelerated decline in lung function in tunnel workers, 663
respiratory complaints, Prognostic factors for respiratory sickness absence and return to work among blue collar workers and office personnel, 246
respiratory effects, Short term exposure to airborne microbial agents during farm work: exposure-response relations with eye and respiratory symptoms, 113
respiratory health, An epidemiological study of the respiratory health of workers in the European refractory ceramic fibre industry, 800
respiratory morbidity, Respiratory health effects from exposure to carbon black: results of the phase 2 and 3 cross sectional studies in the European carbon black manufacturing industry, 496
respiratory symptoms, Experimental exposure to propylene glycol mist in aviation emergency training: acute ocular and respiratory effects, 649
return to work, Prognostic factors for respiratory sickness absence and return to work among blue collar workers and office personnel, 246
risk assessment, Benzene in the environment: an assessment of the potential risks to the health of the population: *review*, 2
Consultative team to assess manual handling and reduce the risk of occupational injury, 339
risk factors, Are occupational, hobby, or lifestyle exposures associated with Philadelphia chromosome positive chronic myeloid leukaemia?, 722
Interrelations of risk factors and low back pain in scaffolders, 597
routine data, Use of routinely collected data on trihalomethane in drinking water for epidemiological purposes, 447

selenium, Cross sectional and longitudinal study on selenium, glutathione peroxidase, smoking, and occupational exposure in coal miners, 239
severity, Occupational injuries in Italy: risk factors and long term trend (1951–98), 330
sewage, Hepatitis A in workers exposed to sewage: a systematic review: *review*, 762
sex, Physical, psychosocial, and organisational factors relative to sickness absence: a study based on Sweden Post, 178

sex ratio, Sex ratio in offspring of those affected by dioxin and dioxin-like compounds: the Yusho, Seveso, and Yucheng incidents, **540**
sheep dip, Estimation of cumulative exposure to organophosphate sheep dips in a study of chronic neurological health effects among United Kingdom sheep dippers, **694**
sheep dippers, An epidemiological study of the relations between exposure to organophosphate pesticides and indices of chronic peripheral neuropathy and neuropsychological abnormalities in sheep farmers and dippers, **702**
shift work, Changes in frequency of premature complexes and heart rate variability related to shift work, **678**
Health effects of shift work and extended hours of work, **68**
Is there an association between shift work and having a metabolic syndrome? Results from a population based study of 27 485 people, **747**
sickness absence, Prognostic factors for respiratory sickness absence and return to work among blue collar workers and office personnel, **246**
Sickness absence in diabetic employees, **432**
The management of sickness absence, **420**
The role of extended weekends in sickness absenteeism, **818**
silicosis, Exposure to silica and silicosis among tin miners in China: exposure-response analyses and risk assessment, **31**
Mortality from lung cancer among silicotic patients in Sardinia: an update study with 10 more years of follow up, **786**
Radiological progression and its predictive risk factors in silicosis, **467**
skin prick test, Role of skin prick test and serological measurement of specific IgE in the diagnosis of occupational asthma resulting from exposure to vinyl sulphone reactive dyes, **411**
sleep reduction, Comparing performance on a simulated 12 hour shift rotation in young and older subjects, **58**
smoking, Dust exposure and impairment of lung function at a small iron foundry in a rapidly developing country, **656**
Relation between exposure to asbestos and smoking jointly and the risk of lung cancer: *review,* **145**
smoking habits, Incidence of cancer among bookbinders, printers, photoengravers, and typesetters, **523**
social conformity, Exposure to organic solvents and personality, **14**
social support, Psychological factors and visual fatigue in working with video display terminals, **267**
solvents, Case-control study of leatherwork and male infertility, **217**
Evaluation of a modified German version of the Q16 questionnaire for neurotoxic symptoms in workers exposed to solvents, **19**
Occupational exposure to solvents and male infertility, **635**
Routine diagnostic procedures for chronic encephalopathy induced by solvents: survey of experts, **382**
spatial random field model, Small area estimation of incidence of cancer around a known source of exposure with fine resolution data, **315**
sperm density, Are problems with male reproductive health caused by endocrine disruption?, **281**
spirometry, Respiratory health effects from exposure to carbon black: results of the phase 2 and 3 cross sectional studies in the European carbon black manufacturing industry, **496**
sports, A prospective study of work related factors and physical exercise as predictors of shoulder pain, **528**
statistics, Reporting of occupational and environmental research: use and misuse of statistical and epidemiological methods: *correspondence,* **63**
strategies, Research priorities in occupational health in Italy, **325**
stress, Association between job strain and prevalence of hypertension: a cross sectional analysis in a Japanese working population with a wide range of occupations: the Jichi Medical School cohort study, **367**
Psychological factors and visual fatigue in working with video display terminals, **267**
styrene, Neurobehavioural changes and persistence of complaints in workers exposed to styrene in a polyester boat building plant: influence of exposure characteristics and microsomal epoxide hydrolase phenotype, **103**
Serum hepatic biochemical activity in two populations of workers exposed to styrene, **95**
surveillance, Serum hepatic biochemical activity in two populations of workers exposed to styrene, **95**
surveillance system, Hospital admissions among male drivers in Denmark, **253**
survey method, Effects of drop out in a longitudinal study of musculoskeletal disorders, **194**
symptoms, Health and exposures of United Kingdom Gulf war veterans. Part I: The pattern and extent of ill health, **291**
New challenges facing ill health in Gulf war veterans: *editorial,* **289**
systematic review, Hepatitis A in workers exposed to sewage: a systematic review: *review,* **762**

TCDD, Dioxin and diabetes mellitus: an analysis of the combined NIOSH and Ranch Hand data, **641**
teams, Consultative team to assess manual handling and reduce the risk of occupational injury, **339**
tear film stability break up time, Experimental exposure to propylene glycol mist in aviation emergency training: acute ocular and respiratory effects, **649**
temperature threshold, Quantitative thermal perception thresholds relative to exposure to vibration, **472**
testicular cancer, Are problems with male reproductive health caused by endocrine disruption?, **281**
thyroid, Association between serum concentrations of hexachlorobenzene and polychlorobiphenyls with thyroid hormone and liver enzymes in a sample of the general population, **172**
time to pregnancy, Solvent use and time to pregnancy among female personnel in biomedical laboratories in Sweden, **225**
toxicity, Ultrafine particles, **211**
traffic fumes, Exposure to benzene in urban workers: environmental and biological monitoring of traffic police in Rome, **165**
traffic police, Exposure to benzene in urban workers: environmental and biological monitoring of traffic police in Rome, **165**
trends, Monitoring occupational diseases: response: *editorial,* **692**
Monitoring trends in occupational illness: *editorial,* **691**
trihalomethanes, Relation between trihalomethane compounds and birth defects, **443**
Use of routinely collected data on trihalomethane in drinking water for epidemiological purposes, **447**
tungsten, Exposure assessment in the hard metal manufacturing industry with special regard to tungsten and its compounds, **631**
type 2 diabetes mellitus, Hours of work and the risk of developing impaired fasting glucose or type 2 diabetes mellitus in Japanese male office workers, **569**

ultrafine particles, Acute respiratory effects of particles: mass or number?, **154**
Ultrafine particles and nitrogen oxides generated by gas and electric cooking, **511**
Ultrafine particles, **211**
unemployment, Mortality among women and men relative to unemployment, part time work, overtime work, and extra work: a study based on data from the Swedish twin registry, **52**

ventilation, Exposure assessment and lung function in pig and poultry farmers, **405**
Functional variables associated with the clinical grade of dyspnoea in coal miners with pneumoconiosis and mild bronchial obstruction, **794**
vibration, Quantitative thermal perception thresholds relative to exposure to vibration, **472**
Raynaud's phenomenon in workers exposed to vibration: *letter,* **279**
vibration frequency, duration, and magnitude, Response of finger circulation to energy equivalent combinations of magnitude and duration of vibration, **185**

water chlorination, Foetal growth and duration of gestation relative to water chlorination, **437**
water pollutants, Public health response to an incident of secondary chemical contamination at a beach in the United Kingdom, **232**
work, Association between job strain and prevalence of hypertension: a cross sectional analysis in a Japanese working population with a wide range of occupations: the Jichi Medical School cohort study, **367**
work environment, Physical, psychosocial, and organisational factors relative to sickness absence: a study based on Sweden Post, **178**
work related symptoms, Short term exposure to airborne microbial agents during farm work: exposure-response relations with eye and respiratory symptoms, **113**
workers, Effects on the nervous system in different groups of workers exposed to aluminium, **453**
working environments, Environmental tobacco smoke and lung function in employees who never smoked: the Scottish MONICA study, **563**

***x* ray fluorescence,** Predictors of DMSA chelatable lead, tibial lead, and blood lead in 802 Korean lead workers, **73**